

# A proof of the equivalence of two complex structures on the punctured tangent bundle of complex projective space

II Kiyotaka<sup>1</sup> and NAKANO Makoto<sup>2</sup>

(Received November 15, 2000)

## Abstract

We give a proof of the equivalence of two complex structures on the punctured tangent bundle  $\mathring{TP}^n(\mathbb{C})$  of complex projective space  $P^n(\mathbb{C})$ .

## 1 Introduction

K. Furutani, R. Tanaka, and S. Yoshizawa constructed a complex (Kähler) structure on the punctured (co-)tangent bundle  $\mathring{TP}^n(\mathbb{C})$  of the complex projective space  $P^n(\mathbb{C})$  by constructing a diffeomorphism of  $\mathring{TP}^n(\mathbb{C})$  onto a “complex cone”  $\mathcal{A}(\mathbb{C})$  in the space  $M(n+1, \mathbb{C})$  of complex matrices ([FT], [FY]). Motivated by their works, we also constructed a complex (Kähler) structure on  $\mathring{TP}^n(\mathbb{C})$  by a Riemannian geometric method ([IY], [Ii], [IM]). Since our complex structure has similar properties as that of Furutani et al., it seems that they coincide, but no proof has been given. The purpose of the present note is to prove that these structures coincide. (cf. [Na])

## 2 Tangent bundle of Riemannian manifold

Let  $M$  be a Riemannian manifold of dimension  $n$  with a Riemannian metric  $g$ . Let  $T_p M$  denote the tangent space to  $M$  at a point  $p$  of  $M$ ,  $TM$  the tangent bundle of  $M$ , and  $\pi$  the bundle projection of  $TM$  onto  $M$ . Let  $T_u TM$  denote the tangent space to  $TM$  at  $u \in TM$ , and  $TTM$  the tangent bundle of  $TM$ . Let  $\nabla$  denote the Levi-Civita connection of  $M$ , and  $\mathcal{K} : TTM \rightarrow TM$  the connection map corresponding to  $\nabla$ . For each  $u \in TM$ ,

---

<sup>1</sup>Division of Mathematical Structures, Department of Mathematical Sciences, Faculty of Science, Yamagata University, Yamagata 990-8560, Japan (e-mail address ii@sci.kj.yamagata-u.ac.jp)

<sup>2</sup>Izumimati 2-3-9, Tendo 994-0015, Japan

let  $T_u^H TM$  (resp.  $T_u^V TM$ ) denote the kernel of  $\mathcal{K}|_{T_u TM}$  (resp.  $d\pi|_{T_u TM}$ ), which is an  $n$ -dimensional subspace of  $T_u TM$  called the horizontal (resp. vertical) subspace of  $T_u TM$ . We have a direct-sum decomposition:

$$T_u TM = T_u^H TM \oplus T_u^V TM.$$

Elements of  $T_u^H TM$  (resp.  $T_u^V TM$ ) are called horizontal (resp. vertical) vectors at  $u$ . If  $u, v \in T_p M$ ,  $v_u^H$  (resp.  $v_u^V$ ) will denote the horizontal (resp. vertical) vector obtained by the horizontal (resp. vertical) lift of  $v$  to  $T_u TM$ . Note that  $v_u^H$  is characterized by

$$\mathcal{K}(v_u^H) = 0 \quad \text{and} \quad d\pi(v_u^H) = v,$$

and  $v_u^V$  is characterized by

$$\mathcal{K}(v_u^V) = v \quad \text{and} \quad d\pi(v_u^V) = 0.$$

Note that the standard almost complex structure  $J_0$  on  $TM$  is a  $(1, 1)$ -tensor field on  $TM$  characterized by

$$J_0(v_u^H) = v_u^V \quad \text{and} \quad J_0(v_u^V) = -v_u^H.$$

(cf. [Do], [GKM], [Sa1], [Sa2])

### 3 Complex projective space

Let  $M(n+1, \mathbb{C})$  denote the space of  $(n+1) \times (n+1)$  complex matrices with the bracket operation  $[A, B] = AB - BA$ , and the norm  $\|A\| = \sqrt{\text{tr}(A^* A)}$ . Let  $Herm(n+1)$  denote the subspace of  $M(n+1, \mathbb{C})$ , consisting of Hermitian matrices with the Euclidean inner product  $(A, B) \mapsto \text{tr}(AB)$ . A vector tangent to  $Herm(n+1)$  at a point  $P \in Herm(n+1)$  is denoted by a pair  $(P, V)$  with  $V \in Herm(n+1)$ . The tangent space to  $Herm(n+1)$  at  $P$  is denoted by

$$T_P Herm(n+1) = \{(P, V) \mid V \in Herm(n+1)\}.$$

The complex projective space  $P^n(\mathbb{C})$  of complex dimension  $n$  is represented as a submanifold of  $Herm(n+1)$  as follows:

$$P^n(\mathbb{C}) = \{P \in Herm(n+1) \mid P^2 = P, \text{tr} P = 1\}.$$

The tangent space to  $P^n(\mathbb{C})$  at a point  $P \in P^n(\mathbb{C})$  is given by

$$T_P P^n(\mathbb{C}) = \{(P, Q) \mid Q \in Herm(n+1), PQ + QP = Q\}.$$

Let  $g$  denote the induced Riemannian metric on  $P^n(\mathbb{C})$ , i.e.,  $g((P, Q), (P, R)) = \text{tr}(QR)$ .

The tangent bundle to  $P^n(\mathbb{C})$  is denoted by  $TP^n(\mathbb{C})$ .

**Lemma 3.1** *The standard complex structure  $j$  on  $P^n(\mathbb{C})$  is given by*

$$j : TP^n(\mathbb{C}) \rightarrow TP^n(\mathbb{C}), \quad j((P, Q)) = (P, \sqrt{-1}[P, Q]).$$

Let  $\mathbf{C}^{n+1}$  be the complex  $(n+1)$ -space with the standard norm  $\| \cdot \|$ ;

$$\mathbf{C}^{n+1} = \{ p = {}^t(p_1, p_2, \dots, p_{n+1}) \mid p_i \in \mathbf{C} \}.$$

**Lemma 3.2** ([FT], [FY]) *For any point  $P$  of  $P^n(\mathbf{C})$ , and for any vector  $(P, Q)$  tangent to  $P^n(\mathbf{C})$  at  $P$ , there exist  $p, q \in \mathbf{C}^{n+1}$  which satisfy*

$$(1) \|p\| = 1, \quad (2) p^*q = 0, \quad (3) P = pp^*, \quad (4) Q = pq^* + qp^*.$$

Note that  $\|Q\|^2 = 2\|q\|^2$ .

**Lemma 3.3** ([FT], [FY]) *Let  $(P, Q)$  and  $(P, R)$  be vectors tangent to  $P^n(\mathbf{C})$  at  $P$ . Then we have*

$$\begin{aligned} (1) \operatorname{tr} Q &= \operatorname{tr}(PQ) = 0, & (2) PQP &= 0, \\ (3) PQ^2 &= Q^2P = \frac{1}{2}\|Q\|^2P, & (4) Q^2 &= \frac{1}{2}\|Q\|^2P + QPQ, \\ (5) Q^3 &= \frac{1}{2}\|Q\|^2Q, & (6) \|PQ\|^2 &= \frac{1}{2}\|Q\|^2, \\ (7) \operatorname{tr}(PQ^2) &= \frac{1}{2}\|Q\|^2, & (8) PQR &= QRP, \\ (9) P(QR + RQ) &= \operatorname{tr}(QR)P, & (10) Q(I - 2P) &= -(I - 2P)Q, \\ (11) \operatorname{tr}([P, Q]Q) &= 0, & (12) Q[P, Q] &= -[P, Q]Q, \\ (13) \|Q\|^2R + 2(I - 2P)[Q^2, R] &= \operatorname{tr}(QR)Q - \operatorname{tr}([P, Q]R)[P, Q], \end{aligned}$$

where  $I$  denotes the unit matrix.

$T_P P^n(\mathbf{C})$  is a linear subspace of the Euclidean space  $T_P \operatorname{Herm}(n+1)$ .

**Lemma 3.4** *The orthogonal projection  $\tau_P : T_P \operatorname{Herm}(n+1) \rightarrow T_P P^n(\mathbf{C})$  is given by*

$$\tau_P(P, V) = (P, PV + VP - 2PVP) = (P, [P, [P, V]]).$$

**Proof**  $(P, PV + VP - 2PVP)$  is tangent to  $P^n(\mathbf{C})$  at  $P$ , since  $(PV + VP - 2PVP)^* = PV + VP - 2PVP$ , and  $P(PV + VP - 2PVP) + (PV + VP - 2PVP)P = PV + VP - 2PVP$ .  $(P, V) - (P, PV + VP - 2PVP) = (P, V - (PV + VP - 2PVP))$  is orthogonal to each vector  $(P, Q)$  tangent to  $P^n(\mathbf{C})$ , since  $\operatorname{tr}(Q(V - (PV + VP - 2PVP))) = 0$ .  $\blacksquare$

Let  $t \mapsto P(t)$  be a  $C^\infty$  curve on  $P^n(\mathbf{C})$ , and  $\xi : t \mapsto \xi(t) = (P(t), Q(t))$  be a vector field along this curve such that  $\xi(t)$  is tangent to  $P^n(\mathbf{C})$  at  $P(t)$ . Then the covariant derivative  $\nabla_{\frac{d}{dt}} \xi : t \mapsto \nabla_{\frac{d}{dt}} \xi(t)$  of  $\xi$  is defined by

$$\nabla_{\frac{d}{dt}} \xi(t) := \tau_{P(t)}(P(t), Q'(t)),$$

which is a vector field along this curve.  $\xi$  is said to be *parallel* if  $\nabla_{\frac{d}{dt}} \xi(t) = (P(t), 0)$  for all  $t$ .

**Lemma 3.5** *Let  $(P, Q), (P, R)$  be vectors tangent to  $P^n(\mathbf{C})$  at  $P$ , and  $t \mapsto P(t)$  be a  $C^\infty$  curve on  $P^n(\mathbf{C})$  such that  $P(0) = P$  and  $\dot{P}(0) := (P(0), P'(0)) = (P, R)$ . Let  $\xi : t \mapsto \xi(t) = (P(t), Q(t))$  be the parallel vector field along this curve with initial condition  $\xi(0) = (P(0), Q(0)) = (P, Q)$ . Then we have*

$$Q'(0) = (I - 2P)(QR + RQ).$$

**Proof** Since  $\xi$  is parallel, we have by Lemma 3.4

$$(P(t), 0) = \nabla_{\frac{d}{dt}} \xi(t) = (P(t), P(t)Q'(t) + Q'(t)P(t) - 2P(t)Q'(t)P(t)).$$

Putting  $t = 0$ , we have

$$PQ'(0) + Q'(0)P - 2PQ'(0)P = 0. \quad (1)$$

Multiplying (1) by  $P$  on the left, we have (since  $P^2 = P$ )

$$PQ'(0) + PQ'(0)P - 2PQ'(0)P = 0. \quad (2)$$

Multiplying (1) by  $P$  on the right, we have

$$PQ'(0)P + Q'(0)P - 2PQ'(0)P = 0. \quad (3)$$

From (2) and (3) we have

$$PQ'(0) = Q'(0)P = PQ'(0)P. \quad (4)$$

On the other hand, since  $\xi(t)$  is tangent to  $P^n(\mathbf{C})$  at  $P(t)$ , we have  $P(t)Q(t) + Q(t)P(t) = Q(t)$  for all  $t$ . Differentiating this with respect to  $t$ , and putting  $t = 0$ , we have  $P(0)Q'(0) + P'(0)Q(0) + Q(0)P'(0) + Q'(0)P(0) = Q'(0)$ . Since  $P(0) = P$ ,  $Q(0) = Q$ , and  $P'(0) = R$ , we have

$$Q'(0) = PQ'(0) + RQ + QR + Q'(0)P. \quad (5)$$

Multiplying (5) by  $P$  on the left, we have  $PQ'(0) = PQ'(0) + PRQ + PQR + PQ'(0)P$ .

Hence

$$PQ'(0)P = -PQR - PRQ. \quad (6)$$

From (4), (5), and (6) we obtain

$$Q'(0) = QR + RQ - 2PQR - 2PRQ = (I - 2P)(QR + RQ). \quad \blacksquare$$

A vector tangent to  $TP^n(\mathbf{C})$  at  $(P, Q)$  is denoted by  $((P, Q), (R, V))$ . The tangent space to  $TP^n(\mathbf{C})$  at a point  $(P, Q)$  is given by

$$T_{(P,Q)}TP^n(\mathbf{C}) = \{((P, Q), (R, V)) \mid R, V \in \text{Herm}(n+1), PR + RP = R, PV + VP + QR + RQ = V\}.$$

The tangent bundle of  $TP^n(\mathbf{C})$  is denoted by  $TTP^n(\mathbf{C})$ . Let  $(P, Q), (P, R) \in T_P P^n(\mathbf{C})$ . Let  $t \mapsto P(t)$  be a  $C^\infty$  curve on  $P^n(\mathbf{C})$  such that  $P(0) = P$  and  $\dot{P}(0) := (P(0), P'(0)) = (P, R)$ , and  $\xi : t \mapsto \xi(t) = (P(t), Q(t))$  be the parallel vector field along this curve with

initial condition  $\xi(0) = (P(0), Q(0)) = (P, Q)$ . Then the horizontal lift  $(P, R)_{(P, Q)}^H$  of  $(P, R)$  to  $T_{(P, Q)}TP^n(\mathbb{C})$  is given by

$$(P, R)_{(P, Q)}^H = \dot{\xi}(0) := (\xi(0), \xi'(0)) = ((P(0), Q(0)), (P'(0), Q'(0))).$$

By Lemma 3.5, we have

$$(P, R)_{(P, Q)}^H = ((P, Q), (R, (I - 2P)(QR + RQ))).$$

The vertical lift  $(P, R)_{(P, Q)}^V$  of  $(P, R)$  to  $T_{(P, Q)}TP^n(\mathbb{C})$  is given by

$$(P, R)_{(P, Q)}^V = \dot{\eta}(0) := (\eta(0), \eta'(0)) = ((P, Q), (0, R)),$$

where  $\eta$  is a curve on  $TP^n(\mathbb{C})$  given by  $\eta(t) = (P, Q + tR)$ . The horizontal subspace and the vertical subspace of  $T_{(P, Q)}TP^n(\mathbb{C})$  are given, respectively, by

$$T_{(P, Q)}^H TP^n(\mathbb{C}) = \{((P, Q), (R, (I - 2P)(QR + RQ))) \mid (P, R) \in T_P P^n(\mathbb{C})\},$$

and

$$T_{(P, Q)}^V TP^n(\mathbb{C}) = \{((P, Q), (0, R)) \mid (P, R) \in T_P P^n(\mathbb{C})\}.$$

The differential  $d\pi$  of the projection  $\pi : TP^n(\mathbb{C}) \rightarrow P^n(\mathbb{C})$ ,  $(P, Q) \mapsto P$  is given by

$$d\pi : TTP^n(\mathbb{C}) \rightarrow TP^n(\mathbb{C}), \quad d\pi((P, Q), (R, V)) = (P, R).$$

The connection map  $\mathcal{K}$  is given by

$$\mathcal{K} : TTP^n(\mathbb{C}) \rightarrow TP^n(\mathbb{C}), \quad \mathcal{K}((P, Q), (R, V)) = (P, PV + VP - 2PVP).$$

Note that

$$\begin{aligned} (1) \quad d\pi((P, R)_{(P, Q)}^H) &= (P, R), & (2) \quad d\pi((P, R)_{(P, Q)}^V) &= (P, 0), \\ (3) \quad \mathcal{K}((P, R)_{(P, Q)}^H) &= (P, 0), & (4) \quad \mathcal{K}((P, R)_{(P, Q)}^V) &= (P, R). \end{aligned}$$

**Lemma 3.6** *Let  $((P, Q), (R, V)) \in T_{(P, Q)}TP^n(\mathbb{C})$ . Then it is represented as a sum of horizontal and vertical vectors as follows:*

$$\begin{aligned} ((P, Q), (R, V)) &= ((P, Q), (R, (I - 2P)(QR + RQ))) \\ &\quad + ((P, Q), (0, V - (I - 2P)(QR + RQ))) \\ &= (P, R)_{(P, Q)}^H + (P, V - (I - 2P)(QR + RQ))_{(P, Q)}^V. \end{aligned}$$

**Proof** It suffices to show that  $(P, V - (I - 2P)(QR + RQ))$  is tangent to  $P^n(\mathbb{C})$  at  $P$ , which follows easily from (8) of Lemma 3.3.  $\blacksquare$

## 4 Complex structure on $\overset{\circ}{T}P^n(\mathbb{C})$

Let  $\overset{\circ}{T}P^n(\mathbb{C})$  denote the punctured tangent bundle of  $P^n(\mathbb{C})$ , i.e.

$$\overset{\circ}{T}P^n(\mathbb{C}) = \{(P, Q) \in TP^n(\mathbb{C}) \mid Q \neq 0\},$$

which is an open subset of  $TP^n(\mathbb{C})$ . We define an almost complex structure  $J$  on  $\overset{\circ}{T}P^n(\mathbb{C})$  by

$$\begin{aligned}
 J((j(P, Q))_{(P, Q)}^H) &= \sqrt{2}\|Q\|(j(P, Q))_{(P, Q)}^V, \\
 J((j(P, Q))_{(P, Q)}^V) &= -\frac{1}{\sqrt{2}\|Q\|}(j(P, Q))_{(P, Q)}^H, \\
 J((P, R)_{(P, Q)}^H) &= \frac{\|Q\|}{\sqrt{2}}(P, R)_{(P, Q)}^V, \\
 J((P, R)_{(P, Q)}^V) &= -\frac{\sqrt{2}}{\|Q\|}(P, R)_{(P, Q)}^H
 \end{aligned}$$

for  $(P, R)$  orthogonal to  $j(P, Q)$ ,  $((P, Q) \in \mathring{T}P^n(\mathbf{C}))$ . It is known that  $J$  is integrable, i.e.,  $J$  is a complex structure on  $\mathring{T}P^n(\mathbf{C})$ . (See [Ii], [IY], [IM].)

**Definition 4.1** ([FT], [FY]) We define a complex submanifold  $\mathcal{A}(\mathbf{C})$  of  $M(n+1, \mathbf{C})$ , and a mapping  $\Phi$  by

$$\begin{aligned}
 \mathcal{A}(\mathbf{C}) &= \{A \in M(n+1, \mathbf{C}) \mid A^2 = 0, \text{rank} A = 1\}, \\
 \Phi : \mathring{T}P^n(\mathbf{C}) &\rightarrow \mathcal{A}(\mathbf{C}), \quad \Phi(P, Q) = \|Q\|^2 P - Q^2 + \frac{\sqrt{-1}}{\sqrt{2}}\|Q\|Q.
 \end{aligned}$$

**Remark** ([FT], [FY]) If we choose  $p, q$  as in Lemma 3.2, then

$$\Phi(P, Q) = (\|q\|p + \sqrt{-1}q)(\|q\|p - \sqrt{-1}q)^*.$$

**Lemma 4.2** ([FT], [FY])  $\Phi$  is well-defined, and is a diffeomorphism.

**Lemma 4.3** Let  $(P, Q) \in \mathring{T}P^n(\mathbf{C})$ . Then we have

(1) For a tangent vector  $((P, Q), (R, V))$  to  $\mathring{T}P^n(\mathbf{C})$  at  $(P, Q)$ ,

$$\begin{aligned}
 d\Phi((P, Q), (R, V)) &= \left( \Phi(P, Q), 2\text{tr}(QV)P + \|Q\|^2 R - (QV + VQ) \right. \\
 &\quad \left. + \frac{\sqrt{-1}}{\sqrt{2}\|Q\|}\text{tr}(QV)Q + \frac{\sqrt{-1}}{\sqrt{2}}\|Q\|V \right).
 \end{aligned}$$

(2) For the horizontal lift to  $(P, Q)$  of a tangent vector  $(P, R)$ ,

$$\begin{aligned}
 d\Phi((P, R)_{(P, Q)}^H) &= \left( \Phi(P, Q), \frac{1}{2}\text{tr}(QR)Q - \frac{1}{2}\text{tr}([P, Q]R)[P, Q] + \frac{1}{2}\|Q\|^2 R \right. \\
 &\quad \left. - \sqrt{-1}\sqrt{2}\|Q\|\text{tr}(QR)P + \frac{\sqrt{-1}}{\sqrt{2}}\|Q\|(QR + RQ) \right).
 \end{aligned}$$

(3) For the vertical lift to  $(P, Q)$  of a tangent vector  $(P, R)$ ,

$$\begin{aligned}
 d\Phi((P, R)_{(P, Q)}^V) &= \left( \Phi(P, Q), 2\text{tr}(QR)P - (QR + RQ) \right. \\
 &\quad \left. + \frac{\sqrt{-1}}{\sqrt{2}\|Q\|}\text{tr}(QR)Q + \frac{\sqrt{-1}}{\sqrt{2}}\|Q\|R \right).
 \end{aligned}$$

(4) For the horizontal lift to  $(P, Q)$  of  $j(P, Q)$ ,

$$d\Phi((j(P, Q))_{(P, Q)}^H) = \left( \Phi(P, Q), \sqrt{-1}\|Q\|^2[P, Q] \right).$$

(5) For the vertical lift to  $(P, Q)$  of  $j(P, Q)$ ,

$$d\Phi((j(P, Q))_{(P, Q)}^V) = \left( \Phi(P, Q), -\frac{1}{\sqrt{2}}\|Q\|[P, Q] \right).$$

**Lemma 4.4** Let  $(P, Q) \in \mathring{TP}^n(\mathbb{C})$ , and  $(P, R)$  be orthogonal to  $j(P, Q)$ . Then we have

$$(1) \quad d\Phi(J((j(P, Q))_{(P, Q)}^H)) = \sqrt{-1} d\Phi((j(P, Q))_{(P, Q)}^H),$$

$$(2) \quad d\Phi(J((j(P, Q))_{(P, Q)}^V)) = \sqrt{-1} d\Phi((j(P, Q))_{(P, Q)}^V),$$

$$(3) \quad d\Phi(J((P, R)_{(P, Q)}^H)) = \sqrt{-1} d\Phi((P, R)_{(P, Q)}^H),$$

$$(4) \quad d\Phi(J((P, R)_{(P, Q)}^V)) = \sqrt{-1} d\Phi((P, R)_{(P, Q)}^V).$$

**Proposition 4.5**  $d\Phi \circ J = \sqrt{-1} d\Phi.$

**Theorem 4.6** If we identify  $\mathring{TP}^n(\mathbb{C})$  with  $\mathcal{A}(\mathbb{C})$  by  $\Phi$ , then the complex structure  $J$  on  $\mathring{TP}^n(\mathbb{C})$  coincides with the complex structure  $\sqrt{-1}$  on  $\mathcal{A}(\mathbb{C})$ .

**Remark** A similar result is obtained for the case of quaternion projective space. (cf. [Ka])

## References

- [DS] A. S. Dancer and R. Szöke, Symmetric spaces, adapted complex structures and hyperKähler structures, Quart. J. Math. Oxford (2) 48 (1997), 27-38.
- [Do] P. Dombrowski, On the geometry of the tangent bundle, J. Reine Angew. Math. 210 (1962), 73-88.
- [FT] K. Furutani and R. Tanaka, A Kähler structure on the punctured cotangent bundle of complex and quaternion projective spaces and its application to a geometric quantization I, J. Math. Kyoto Univ. 34-4 (1994), 719-737.
- [FY] K. Furutani and S. Yoshizawa, A Kähler structure on the punctured cotangent bundle of complex and quaternion projective spaces and its application to a geometric quantization II, Japanese J. Math. 21 (1995), 355-392.
- [GKM] D. Gromoll, W. Klingenberg and W. Meyer, Riemannsche Geometrie im Großen, Lecture Notes in Math. 55, Springer-Verlag, 1968.
- [Ii] K. Ii, Kähler structures on the tangent bundle of Riemannian manifolds of constant positive curvature, Preprint (1996).
- [IM] K. Ii and T. Morikawa, Kähler structures on the tangent bundle of Riemannian manifolds of constant positive curvature, Bull. Yamagata Univ., Nat. Sci. 14 (1999), 141-154.

- [IY] K. Ii and S. Yaguchi, On a Riemannian geometric construction of a Kähler structure on the punctured tangent bundle of complex projective space, *Bull. Yamagata Univ., Nat. Sci.* 14 (1997), 59-70.
- [Ka] K. Kamata, On the equivalence of complex structures on the tangent bundle of quaternion projective space, Thesis, Yamagata University, 1999.
- [LS] L. Lempert and R. Szőke, Global solutions of the homogeneous complex Monge-Ampère equation and complex structures on the tangent bundles of Riemannian manifolds, *Math. Ann.* 290 (1991), 689-712.
- [MN] A. Morimoto and T. Nagano, On pseudo-conformal transformations of hypersurfaces, *J. Math. Soc. Japan* 15 (1963), 289-300.
- [Na] M. Nakano, On the equivalence of complex structures on the tangent bundle of complex projective space, Thesis, Yamagata University, 1998.
- [Ra1] J. H. Rawnsley, Coherent state and Kaehler manifold, *Quart. J. Math. Oxford* (2) 28 (1977), 403-415.
- [Ra2] J. H. Rawnsley, A nonunitary pairing of polarizations for the Kepler problem, *Trans. Amer. Math. Soc.* 250 (1979), 167-180.
- [Sa1] S. Sasaki, On the differential geometry of tangent bundles of Riemannian manifolds, *Tôhoku Math. J.* 10 (1958), 338-354.
- [Sa2] S. Sasaki, On the differential geometry of tangent bundles of Riemannian manifolds, Part II, *Tôhoku Math. J.* 14 (1962), 146-155.
- [So] J. M. Souriau, *Sur la variété de Kepler*, *Symposia Math.* 14, Academic Press, 1974, 343-360.
- [Sz1] R. Szőke, Adapted complex structures and geometric quantization, *Nagoya Math. J.* 154 (1999), 171-183.
- [Sz2] R. Szőke, Complex structures on tangent bundles of Riemannian manifolds, *Math. Ann.* 291 (1991), 409-428.
- [TW1] M. Tahara and Y. Watanabe, Natural almost Hermitian and Kähler metrics on the tangent bundles, *Math. J. Toyama Univ.* 20 (1997), 149-160.
- [TW2] M. Tahara, L. Vanhecke and Y. Watanabe, New structures on tangent bundles, Preprint (1998).